



Cellulitis in chronic oedema of the lower leg: an international cross-sectional study

E.A. Burian ¹, T. Karlsmark,¹ P.J. Franks,² V. Keeley,³ I. Quéré⁴ and C.J. Moffatt ^{1,5,6}

¹Department of Dermato-Venereology & Wound Healing Centre, Bispebjerg Hospital, Copenhagen, Denmark

²Centre for Research and Implementation of Clinical Practice, London, UK

³Lymphoedema Department, Royal Derby Hospital, Derby and University of Nottingham Medical School, Nottingham, UK

⁴Department of Vascular Medicine, Montpellier University Hospital Centre, University of Montpellier, France

⁵School of Social Sciences, Nottingham Trent University, Nottingham, UK

⁶Nottingham University Hospitals NHS Trust, Nottingham, UK

Linked Comment: P. S. Mortimer. *Br J Dermatol* 2021; **185**:10–11.

Summary

Correspondence

Christine J. Moffatt.

Email: christine.moffatt@ntu.ac.uk

Accepted for publication

5 January 2021

Funding sources

The study was funded by the registered charity International Lymphoedema Framework and the medical device company 3M Healthcare. 3M Healthcare had no role in the study design, data collection, data analysis, data interpretation, writing or decision to submit the article for publication. All authors and researchers acted independently from funders.

Conflicts of interest

See Appendix for full details.

DOI 10.1111/bjd.19803

Background Cellulitis and chronic oedema are common conditions with considerable morbidity. The number of studies designed to assess the epidemiology of cellulitis in chronic oedema is scarce.

Objectives To investigate the prevalence and risk factors of cellulitis in chronic leg oedema, including lymphoedema.

Methods A cross-sectional study included 40 sites in nine countries during 2014–17. Adults with clinically proven unilateral or bilateral chronic oedema (oedema > 3 months) of the lower leg were included. The main outcome measures were frequency and risk factors for cellulitis within the last 12 months.

Results Out of 7477 patients, 15.78% had cellulitis within the last 12 months, with a lifetime prevalence of 37.47%. The following risk factors for cellulitis were identified by multivariable analysis: wounds [odds ratio (OR) 2.37, 95% confidence interval (CI) 2.03–2.78], morbid obesity (OR 1.51, 95% CI 1.27–1.80), obesity (OR 1.21, 95% CI 1.03–1.41), midline swelling (OR 1.32, 95% CI 1.04–1.66), male sex (OR 1.32, 95% CI 1.15–1.52) and diabetes (OR 1.27, 95% CI 1.08–1.49). Controlled swelling was associated with a reduced risk (OR 0.59, 95% CI 0.51–0.67). In a subgroup analysis, the risk increased with the stage of oedema [International Society of Lymphology, stage II OR 2.04 (95% CI 1.23–3.38) and stage III OR 4.88 (95% CI 2.77–8.56)].

Conclusions Cellulitis in chronic leg oedema is a global problem. Several risk factors for cellulitis were identified, of which some are potentially preventable. Our findings suggest that oedema control is one of these. We also identified that advanced stages of oedema, with hard/fibrotic tissue, might be an important clinical indicator to identify patients at particular risk.

What is already known about this topic?

- Chronic oedema has many different causes and is a frequent but neglected health-care problem.
- The association between chronic oedema and cellulitis is known, but few studies have clinically evaluated the size of the problem and the risk factors.
- Guidelines suggest that control of oedema is important to reduce the risk of recurrent cellulitis, but the evidence is limited.

What does this study add?

- Cellulitis in chronic leg oedema is common in all countries and types of health facilities.
- Wounds, obesity, midline swelling, male sex and diabetes were independently associated with a recent episode of cellulitis (within the last 12 months).
- Severe stages of oedema were associated with cellulitis, while controlled swelling was associated with a reduced risk.
- Measures to improve the control of swelling may have a major effect on the incidence of cellulitis, being potentially preventable.

Cellulitis is a common bacterial infection of the dermis and subcutaneous tissue,¹ and can occur in any body site, lower limbs being affected in 70–80% of cases.² It is a common medical emergency, often leading to hospitalization, long-term morbidity and recurrent disease.¹ In 2018–19 cellulitis accounted for 1.4% of all emergency admissions in the UK,³ and it has been reported that it is one of the leading causes of potentially preventable hospitalizations.⁴

Chronic oedema is also frequent, with an estimated prevalence of 38% in European hospitals,⁵ and 57% of patients cared for by community nurses in the UK.⁶ Yet, it is a neglected healthcare issue.⁷ Chronic oedema is defined as oedema present for more than 3 months.⁸ Traditionally, the term lymphoedema has been used for oedema resulting from a failure of the lymphatics, e.g. due to congenital malformation, cancer, injury or filariasis. However, recent research indicates the substantial role of the lymphatics in all chronic oedema, leading to the introduction of this umbrella term. Chronic oedema is often multifactorial and covers a wide range of pathologies including lymphoedema (primary and secondary), but also swelling due to venous insufficiency, immobility and obesity.⁷

A recent meta-analysis identified lymphoedema/chronic leg oedema as an independent risk factor for cellulitis [odds ratio (OR) 6.77, 95% confidence interval (CI) 3.46–13.27].⁹ One of the reasons that the risk of cellulitis may be increased is due to the important role of the lymphatics in immunity.^{10,11} Although both diseases are common, few studies have been designed to clinically examine the epidemiology of cellulitis in patients with chronic oedema. The objective of this study was to investigate the association of potential risk factors with the presence of cellulitis in patients with chronic leg oedema. The identification of preventable or modifiable risk factors could improve patient outcomes.

Methods

This is an international, multicentre, cross-sectional study, performed as part of LIMPRINT (Lymphoedema IMPact and PRevalence – INternational Lymphoedema Framework), an epidemiology study designed to prospectively determine the impact and prevalence of chronic oedema within health services. Forty sites from nine countries participated between June

2014 and August 2017. Hospital (in- and outpatients) and community cases were included. Main outcome of interest was the presence of cellulitis (yes/no) in the sites affected by chronic leg oedema within the last 12 months, and its relation to potential risk factors. Each country and study centre gained the appropriate approvals from the relevant Ethical Review Committee and other research and service development committees.

Chronic oedema

Chronic oedema is defined as oedema present for more than 3 months and affecting one or more areas of the body [limbs, hands/feet, upper body (breast/chest wall, shoulder, back), lower body (buttocks, abdomen), genitalia (scrotum, penis, vulva), head, neck or face].⁸ Oedema was confirmed using the validated 'Pitting Oedema Test' and Stemmer's sign (a positive Stemmer's sign: a skin fold that cannot be pinched at the base of the second toe, which is diagnostic of lymphoedema). Patients with long-standing chronic nonpitting oedema with fibrosis were also included.¹² The duration was determined from medical records and through the patient or caregiver. The severity was judged by palpation and clinical evaluation of the skin, using International Society of Lymphology (ISL) staging,¹³ originally developed for lymphoedema:

Stage I: Early onset, with an accumulation of tissue oedema that decreases with limb elevation; the oedema may be pitting.

Stage II: Limb elevation alone rarely reduces swelling and pitting is manifested.

Stage III: The tissue is fibrotic (hard) and pitting is absent; skin changes such as thickening, hyperpigmentation, increased skin folds, fat deposits and warty overgrowths develop.

Cellulitis

Cellulitis is defined as an acute onset of soft-tissue erythema, warmth and tenderness that rapidly resolves with antibiotics, most often caused by *Streptococcus pyogenes* and/or *Staphylococcus aureus* (to a lesser extent). Erysipelas is a similar infection, but typically affects the more superficial part of the skin compared with cellulitis. The terms are often used interchangeably,¹ and are considered as one clinical issue in this manuscript. The current presence or history of cellulitis were confirmed by a

combination of physical examination, interview with the patient and/or review of the medical records by teams of clinicians, all of which included experts in lymphology.

Study population

The study population comprised adults 18 years of age and older, with clinically proven unilateral or bilateral chronic leg oedema (regardless of the underlying cause), who could understand the study and give informed consent according to ethical standards. Cases were excluded if unwilling or unable to participate, receiving end-of-life care or if judged as not in the patient's best interest.

Data collection

The methods have been published previously.¹⁴ In brief, a standardized core tool was used for all participants; it was developed by an international expert panel and included both a questionnaire and a physical examination. An expert review deemed the tool to be highly accurate.¹⁴ Data were collected by trained healthcare professionals. Lymphoedema specialists confirmed the underlying chronic oedema classification and the diagnosis of cellulitis. An additional tool was used in some centres with the appropriate expertise for undertaking the staging procedure (ISL). All sites followed the international study protocol and complied with standard operating procedures. In nine lymphoedema specialist centres, data using the LIMPRINT core tool were obtained from clinical records of all patients.¹⁵

Variables

There are no internationally agreed definitions on the outcome on chronic oedema management. In this study 'control of swelling' was a subjective judgement by the investigator based on the clinical observation of the limb, clarified with the caregiver and if necessary with the lead physician within each service.¹⁶ It was assessed as either present, absent or 'don't know', at the time of clinical assessment. The type (or absence) of treatment was noted including skin care, exercise, manual lymph drainage, types of compression, antibiotics, psychological support and surgical treatments. Data included demographics and relevant comorbidities. Body mass index (BMI) was estimated according to World Health Organization categories as either underweight (BMI < 20), normal weight (BMI 20–30), obese (BMI 30–40) or morbidly obese (BMI > 40). The site of chronic oedema was collected using a body map where the upper and lower extremities, trunk including genitals (collectively termed midline swelling), face and neck were recorded. The oedema was further classified as either primary (congenital) or secondary (acquired), and whether related to cancer or not. Cancer-related oedema was classified as either caused by treatment and/or due to metastatic disease. Noncancer oedema was classified as due to clinically assessed venous disease (including confirmation by

ultrasound), obesity, immobility, lymphatic filariasis and/or 'other'. Duration of oedema and leg mobility was documented. Wounds defined as 'loss of intact skin' were determined through clinical examination (wound classification will be published elsewhere). In selected centres, the severity of oedema was also assessed using the ISL staging tool.

Statistics

Statistical analyses were performed in Stata 12 (Statacorp; College Station, TX, USA). Due to the explorative study design, a formal sample size determination was not performed. A sample of over 5000 patients was expected to reveal the major factors associated with cellulitis. The principal analysis examined the binary outcome (history vs. no history of cellulitis within the previous 12 months). Factors tested for an association with the outcome were demographics, medical history and leg and swelling characteristics. These variables were chosen as they were believed to be potentially associated with the outcome and could be reliably collected in an international study like this. The principal analysis used logistic regression. Univariate comparisons were followed by a multivariable model, using a stepwise elimination until all factors remaining had an α of < 0.05. Results were presented as ORs and 95% CIs. A similar analysis examined the severity of chronic oedema in a subgroup of 966 patients. Missing data were not imputed and therefore remained missing.

Results

Characteristics of countries, sites and patients

From those included in the Limprint database, 10 127 patients undertook the core questionnaire, had a site of swelling and were 18 years of age or older; 7722 (76.25%) of these were identified with leg oedema and were included in this study. Of these patients with leg oedema, 7477 (96.8%) patients gave a response to the question about whether they had experienced a recent (< 12 months) history of cellulitis. Of the cohort (with or without cellulitis), 61.87% had well-controlled chronic oedema. Patient characteristics are presented in Table 1. In total, 40 sites from nine countries participated, including Australia, Canada, Denmark, France, Ireland, Italy, Japan, Turkey and the UK (Table 2). These included specialist lymphoedema services (73.4%), outpatient acute hospitals (9.0%), hospitalized cases (8.6%), community nursing (1.2%), elderly care residential homes (0.3%), nursing homes (0.1%) and other (7.3%).

General risk factors

On univariate analysis statistically significant associations were found between recent cellulitis and diabetes (OR 1.56), male sex (OR 1.47), morbid obesity (OR 1.56), obesity (OR 1.19), chair-bound patients (OR 1.39), peripheral arterial disease (OR 1.37) and heart failure/ischaemic heart disease (OR 1.25). Age was weakly associated ($P = 0.12$); see Table 3.

Table 1 Demographics of the patients with chronic leg oedema for whom information on presence or absence of cellulitis is available (n = 7477)^a

Characteristic(s)	Number of patients (%) ^b	Missing data
Age, mean years (SD)	65.05 (16.36)	2
Female	5265 (70.42)	0
Weight		11
Normal weight	3120 (41.79)	
Underweight	164 (2.20)	
Obese	2631 (35.24)	
Morbidly obese	1551 (20.77)	
Concomitant disease		
Diabetes	1379 (18.44)	0
Heart failure/ischaemic heart disease	1184 (15.84)	0
Facility		0
Hospital-based cases	7018 (93.86)	
Community cases	125 (1.67)	
Other	334 (4.47)	
Classification of chronic oedema		
Primary	1396 (18.83)	65
Secondary	6016 (81.17)	
Related to cancer or its treatment	1057 (17.63)	22
Noncancer	4937 (82.37)	
Venous disease ^c	2421 (49.12)	
Immobility ^c	1847 (37.47)	
Obesity ^c	1478 (29.99)	
Filariasis ^c	8 (0.22)	
Unilateral leg oedema	1861 (24.89)	0
Bilateral leg oedema	5616 (75.11)	
ISL scale (n = 966) ^d		0
I	237 (24.53)	
II	549 (56.83)	
III	180 (18.63)	
Duration of leg oedema		11
< 1 year	833 (11.16)	
1–2 years	739 (9.90)	
2–5 years	1561 (20.91)	
> 5–10 years	1685 (22.57)	
> 10 years	2648 (35.47)	
Mobility		9
Normal	4203 (56.28)	
Walking aid	2466 (33.02)	
Chair-bound	691 (9.25)	
Bed-bound	108 (1.45)	
Concurrent swelling		
Upper limb	360 (4.81)	0
Midline	586 (7.84)	0
Presence of a leg wound	1129 (15.13)	17
Treatment with compression therapy		20
Compression garment	5101 (68.41)	
Multilayer bandage	1888 (25.32)	
Compression wrap	668 (8.96)	
At least one of the above	5804 (77.83)	
No compression	1653 (22.17)	
Good control of swelling	4314 (61.87)	504 ^e
Life history of cellulitis	2802 (37.47)	
Antibiotics	448 (6.01)	20

(continued)

Table 1 (continued)

Characteristic(s)	Number of patients (%) ^b	Missing data
Hospitalized cases due to cellulitis within the last 12 months	368 (4.95)	47

^aNumbers indicate the actual number of recorded values excluding missing data. ^b%, unless otherwise noted; ^cbreakdown and percentages derived from 'present' values shown in Table 4; ^dISL scale, International Society of Lymphology scale (as described in Patients and Methods) – this was performed only in those centres with the appropriate expertise for undertaking the staging procedure; therefore, missing data is regarded as = 0; ^eIncludes missing and uncertain values.

Local risk factors

On univariate analysis, wounds were identified as a statistically significant risk factor (OR 2.75). Secondary lymphoedema was associated with cellulitis when compared with primary lymphoedema (OR 1.25), but the risk was not related to whether the oedema was caused by cancer or its treatment or a non-cancer cause. Of the other factors only venous disease (OR 1.21) showed a positive association, and concomitant midline swelling (OR 1.30). Control of swelling was associated with a significantly lower risk (OR 0.51) (Table 4).

Independent risk factors

Factors remaining after multivariable analysis (logistic regression) were wounds (OR 2.37), morbid obesity (OR 1.51), obesity (OR 1.21), midline swelling (OR 1.32), male sex (OR 1.32) and diabetes (OR 1.27). Patients with controlled swelling had a markedly lower risk of cellulitis, OR 0.59 (95% CI 0.51–0.67, P < 0.001) (see Table 5).

Frequency of cellulitis

Of the patients with chronic oedema of the lower leg 37.47% (2802 of 7477) experienced at least one episode of cellulitis during their lifetime. In total, 15.78% (n = 1180) had a history of cellulitis within the last 12 months, of whom 368 (31.2%) were hospitalized. The frequency of recent cellulitis ranged from 13.94% in the UK to 38.24% in Canada (see Table 2). The difference is likely explained by the type of facility.

In those assessed for the severity of the chronic oedema (n = 966), the frequency of a recent history of cellulitis changed with ISL stage, affecting 11.56% with ISL stage I, 50.75% in stage II and 37.69% in stage III (Table 6).

Severity of oedema

Severity of oedema (n = 966, Table 6) was significantly associated with cellulitis: ISL stage II OR 2.10 and stage III OR 6.65 compared with stage I, by univariate analysis. An

Table 2 History of lower leg cellulitis (< 12 months) in patients with chronic oedema by country

Country	Total number of patients with chronic oedema	History of cellulitis (< 12 months)	Percentage
UK	4714	657	13.94
France	347	49	14.12
Japan	82	14	17.07
Denmark	859	149	17.35
Italy	1065	211	19.81
Turkey	216	43	19.91
Australia	108	26	24.07
Ireland	18	5	27.78
Canada	68	26	38.24
Total	7477	1180	15.78

increased risk was also seen in hard (fibrotic) tissue vs. soft tissue (OR 2.85), and with a positive Stemmer's sign (OR 2.23). Even after adjustment for sex, obesity, diabetes,

wounds, controlled swelling and midline swelling ($n = 889$, Table 6), ISL stage II yielded an OR of 2.04, and stage III an OR of 4.88, by multivariable analysis.

Discussion

This large study confirms that cellulitis is common in patients with chronic leg oedema. Of the patients studied, 15.78% had experienced at least one episode of cellulitis within the last 12 months, with a lifetime prevalence of 37.47%. The methodology adopted, with a physical examination, access to lymphoedema experts, use of international definitions and standard operating procedures, strengthens the validity of our data. The lifetime prevalence is higher than previously reported (7.95–35.7%), with direct comparison challenging due to methodological differences.^{17–19}

Wounds, obesity, male sex, diabetes, midline swelling and, particularly, advanced stages of chronic oedema were independent risk factors for cellulitis, while control of swelling was associated with a lower risk. Although risk factors in cellulitis have been studied in a meta-analysis (identifying previous

Table 3 Explanatory variables for recent (< 12 months) cellulitis in patients with chronic oedema of the lower leg, by univariate analysis ($n = 7477$)^a

Risk factor	No cellulitis, N (%)	Cellulitis, N (%)	OR (95% CI)	P-value
Sex				
Female	4517 (71.73)	748 (63.39)	1.00	
Male	1780 (28.27)	432 (36.61)	1.47 (1.29–1.66)	< 0.001
Age				
< 45 years	794 (12.61)	140 (11.87)	1.00	
45–64 years	1945 (30.89)	394 (33.42)	1.15 (0.93–1.42)	
65–74 years	1474 (23.41)	289 (24.51)	1.11 (0.89–1.38)	0.12
75–84 years	1389 (22.06)	251 (21.29)	1.02 (0.82–1.28)	
85 + years	694 (11.02)	105 (8.91)	0.86 (0.65–1.13)	
Obesity				
Normal weight	2694 (42.85)	426 (36.13)	1.00	
Underweight	136 (2.16)	28 (2.37)	1.30 (0.86–1.98)	
Obese	2213 (35.20)	418 (35.45)	1.19 (1.03–1.38)	
Morbidly obese	1244 (19.79)	307 (26.04)	1.56 (1.33–1.83)	< 0.001
Leg mobility				
Walks unaided	3570 (56.77)	633 (53.64)	1.00	
Walks with aid	2068 (32.89)	398 (33.73)	1.09 (0.95–1.24)	
Chair-bound	554 (8.81)	137 (11.61)	1.39 (1.14–1.71)	0.007
Bed-bound	96 (1.53)	12 (1.02)	0.70 (0.38–1.29)	
Diabetes				
Absent	5208 (82.71)	890 (75.42)	1.00	
Present	1089 (17.29)	290 (24.58)	1.56 (1.34–1.81)	< 0.001
Heart failure/ischaemic heart disease				
Absent	5332 (84.68)	963 (81.61)	1.00	
Present	965 (15.32)	217 (18.39)	1.25 (1.06–1.46)	0.008
Neurological disease				
Absent	5729 (91.17)	1067 (90.81)	1.00	
Present	555 (8.83)	108 (9.19)	1.04 (0.84–1.30)	0.69
Peripheral arterial disease				
Absent	6088 (96.68)	1127 (95.51)	1.00	
Present	209 (3.32)	53 (4.49)	1.37 (1.01–1.86)	0.044

^aNumbers indicate the actual number of recorded values excluding missing data.

Table 4 Explanatory variables for cellulitis in patients with chronic oedema of the lower leg, by univariate analysis (n = 7477)^a

Risk factor	No cellulitis, N (%)	Cellulitis, N (%)	OR (95% CI)	P-value
Swelling duration, n = 7466				
< 1 year	684 (10.88)	149 (12.64)	1.00	
1–2 years	645 (10.26)	94 (7.97)	0.67 (0.51–0.88)	
2–5 years	1337 (21.27)	224 (19.00)	0.77 (0.61–0.96)	0.015
5–10 years	1399 (22.25)	286 (24.26)	0.94 (0.75–1.17)	
> 10 years	2222 (35.34)	426 (36.13)	0.88 (0.72–1.08)	
Classification, n = 7412				
Primary	1208 (19.35)	188 (16.07)	1.00	
Secondary	5034 (80.65)	982 (83.93)	1.25 (1.06–1.48)	0.008
Secondary cause, n = 5994				
Cancer	897 (17.89)	160 (16.34)	1.00	
Noncancer	4118 (82.11)	819 (83.66)	1.11 (0.93–1.34)	0.25
Cancer cause, n = 1053 ^b				
Cancer treatment				
Absent	125 (13.98)	13 (8.18)	1.00	
Present	769 (86.02)	146 (91.82)	1.83 (1.00–3.32)	0.046
Cancer metastasis				
Absent	789 (88.26)	149 (93.71)	1.00	
Present	105 (11.74)	10 (6.29)	0.50 (0.26–0.99)	0.042
Noncancer, n = 4929 ^c				
Venous				
Absent	2125 (51.67)	383 (46.94)	1.00	
Present	1988 (48.33)	433 (53.06)	1.21 (1.04–1.40)	0.014
Immobility				
Absent	2585 (62.85)	497 (60.91)	1.00	
Present	1528 (37.15)	319 (39.09)	1.09 (0.93–1.27)	0.30
Obesity				
Absent	2952 (71.77)	499 (61.15)	1.00	
Present	1161 (28.23)	317 (38.85)	1.62 (1.38–1.89)	< 0.001
Concomitant arm swelling				
Absent	5990 (95.12)	1127 (95.51)	1.00	
Present	307 (4.88)	53 (4.49)	0.92 (0.68–1.24)	0.57
Concomitant midline swelling				
Absent	5824 (92.49)	1067 (90.42)	1.00	
Present	473 (7.51)	113 (9.58)	1.30 (1.05–1.62)	0.015
Leg wound, n = 7460				
Absent	5490 (87.36)	841 (71.51)	1.00	
Present	794 (12.64)	335 (28.49)	2.75 (2.38–3.19)	< 0.001
Control of swelling, n = 6973				
Not controlled	2082 (35.54)	577 (51.80)	1.00	
Controlled	3777 (64.46)	537 (48.20)	0.51 (0.45–0.58)	< 0.001

^aNumbers indicate the actual number of recorded values excluding missing data. ^bEvidence of the type of cancer cause not given, n = 4.

^cFilariasis was not included in this table due to small numbers; also, missing values, n = 8.

cellulitis, concurrent wounds, leg ulcers, excoriating skin diseases, tinea pedis, obesity and lymphoedema/chronic oedema as risk factors),⁹ only one single-centre study has been specifically designed in patients with chronic oedema/lymphoedema.²⁰ Independent risk factors were percentage difference of circumference of the limb, 'food induced complications experiences', systolic blood pressure and primary lymphoedema. In contrast, we found that secondary lymphoedema was associated with cellulitis on univariate analysis, but it was not an independent risk factor.

Our most important findings were that control of swelling was associated with a significantly lower risk of cellulitis (OR

0.59) while advanced stages of chronic oedema were strong risk factors (ISL stage II: OR 2.04 and stage III: OR 4.88), assessed by multivariable analysis, indicating that cellulitis may be preventable. Measures to control the swelling and halt the progression into advanced stages, e.g. with appropriate compression garment, should be mandatory.

Chronic oedema management is already widely recognized as an adjuvant to antibiotic prophylaxis for recurrent cellulitis.^{21,22} A randomized clinical trial (n = 84) in compression therapy significantly lowered the incidence of recurrence of cellulitis compared with conservative treatment, with a relative risk of 0.37 in favour of compression.²³ However, in our

Table 5 Logistic regression analysis. Independent risk factors associated with cellulitis of the lower leg in patients with chronic oedema (n = 6947)

	OR (95% CI)	P-value
Sex		
Female	1.00	
Male	1.32 (1.15–1.52)	< 0.001
Weight		
Normal weight	1.00	
Underweight	1.17 (0.78–1.82)	
Obese	1.21 (1.03–1.41)	
Morbidly obese	1.51 (1.27–1.80)	< 0.001
Diabetes		
Absent	1.00	
Present	1.27 (1.08–1.49)	0.003
Wound		
Absent	1.00	
Present	2.37 (2.03–2.78)	< 0.001
Midline swelling		
Absent	1.00	
Present	1.32 (1.04–1.66)	0.020
Control of oedema		
Not controlled	1.00	
Controlled	0.59 (0.51–0.67)	< 0.001

cohort only 48.2% of those with recent cellulitis had proper oedema control, highlighting the need to focus on this issue. A reduced incidence of cellulitis by implementation of compression therapy has been reported to decrease healthcare costs almost threefold over a 1-year period, mainly explained by a reduction of acute care costs.²⁴ Mechanisms of compression include reduced capillary filtration, increased lymphatic

drainage and a downregulation of proinflammatory cytokines. Warty skin, venous eczema and occasionally fibrosis (especially seen in ISL stage III) can be reversed.^{25,26} These conditions are reservoirs or entry points for microbes.

It is hypothesized that the risk of cellulitis is increased due to a local immune deficiency, with an ineffective transport of antigens to the lymph node. Lymph stasis may also facilitate bacterial growth and impede bacterial and toxin clearance.²⁷ Cellulitis also seems to impair the lymphatics. A single methicillin-resistant *Staphylococcus aureus* infection has been shown to inhibit lymphatic vessel contraction and flow long after infection clearance.²⁷ As oedema predisposes to cellulitis and cellulitis can impair the lymphatics, potential prophylactic interventions should target all steps in this vicious cycle.¹¹

As expected, wounds were associated with cellulitis, as was obesity. Increased fat deposition in primary lymphoid organs leading to alterations of the leucocyte population might play a part.²⁸ Furthermore, decreased lymphatic transport, fewer lymphatic vessels, and changed architecture and smaller lymph nodes have been observed in obese mice.²⁹ We also found that males were 30% more likely to experience cellulitis than females, which has not been reported.⁹ Male predominance may be due to behavioural and biological factors³⁰ with less-efficient antigen presentation, lower phagocytic activity and lower antibody production.³¹

Diabetes as a risk factor for cellulitis could not be confirmed in the previously mentioned meta-analysis,⁹ perhaps due to a too-small sample size. However, our results are supported by a large, matched cohort study [type I diabetes OR 2.84 (95% CI 2.48–3.25); type II OR 2.03 (1.97–2.08)].³² Abnormal neutrophil function and T-lymphocyte responses might come into play.³² Proper foot care to prevent cellulitis is mandatory for all diabetics.

Table 6 Explanatory variables for cellulitis related to the severity of chronic leg oedema, a subgroup analysis (n = 966)

	No cellulitis, n (%)	Cellulitis, n (%)	OR 95%CI	P-value
Pitting				
Nonpitting	215 (28.0)	62 (31.2)	1.00	0.39
pitting	552 (72.0)	137 (68.8)	0.86 (0.61–1.21)	
Tissue quality				
Soft	575 (74.97)	102 (51.3)	1.00	< 0.001
Hard (fibrotic)	192 (25.0)	97 (48.7)	2.85 (2.07–3.93)	
Stemmer's sign (missing, n = 24)				
Negative	315 (42.0)	47 (24.5)	1.00	< 0.001
Positive	435 (58.0)	145 (75.5)	2.23 (1.56–3.20)	
ISL scale				
Stage I	214 (27.9)	23 (11.6)	1.00	< 0.001
Stage II	448 (58.4)	101 (50.8)	2.10 (1.30–3.39)	
Stage III	105 (13.7)	75 (37.7)	6.65 (3.94–11.20)	
ISL scale after adjustment ^a (n = 889)				
Stage I			1.00	< 0.001
Stage II			2.04 (1.23–3.38)	
Stage III			4.88 (2.77–8.56)	

ISL scale, International Society of Lymphology scale (as described in Patients and Methods). ^aISL scale after adjustment for sex, obesity, diabetes, wound, control and midline swelling by logistic regression.

Being chair-bound was also a risk factor, which might be explained by the increased hydrostatic pressure and lack of use of the calf muscles, worsening the oedema. However, this association disappeared when correcting for other factors.

Limitations to our study need to be noted. Cellulitis is known for being easily misdiagnosed (up to 30·7% of cases).³³ Observer bias was minimized by the use of international definitions of cellulitis, standard operating procedures and training.¹⁴ Recall bias was minimized by seeking information from the medical records.

Although recruitment was made from hospitals and community facilities, the majority of the patients were included from hospitals, potentially skewing our data towards more severe cases. Also, patients from developing countries were not included.

The assessment of 'control of swelling' was open to personal interpretation. However, data were excluded when the clinician was uncertain of the control status ($n = 465$) and access to lymphoedema teams assisted in increasing the accuracy of the decisions.

The lack of correction for prophylactic antibiotics might also influence our data. Although antibiotic use was recorded in 194 of 6283 (3·09%) with no cellulitis vs. 254 of 1174 (21·64%) with a recent history of cellulitis, this did not specify its use in prophylaxis, and may be the result of treating the acute infection. Lastly, one should keep in mind that a cross-sectional study is limited to an assessment at only one timepoint, indicating association but not causation.

In conclusion, our findings confirm that cellulitis in chronic leg oedema is a global problem. Although guidelines support the use of oedema control to prevent cellulitis (e.g. with compression therapy), a substantial number of those recruited had uncontrolled swelling. This study adds epidemiological evidence of what has been known anecdotally for a long time: that oedema control is associated with a lower risk of cellulitis. Wounds, warty skin and eczema, as often seen in advanced stages of oedema, are potential entry points and/or reservoirs for microbes, and can be prevented and treated by compression therapy. Prevention of deterioration of the oedema may have a significant effect on reducing the risk of cellulitis, and thereby reducing healthcare costs.

Acknowledgments

3 M Healthcare provided the International Lymphoedema Framework an unrestricted research grant for the LIMPRINT project and access and support of an electronic case report form. We are grateful to all the patients and staff participating in this study.

References

- Raff AB, Kroshinsky D. Cellulitis: a review. *JAMA* 2016; **316**:325–37.
- Hirschmann JV, Raugi GJ. Lower limb cellulitis and its mimics: part I. Lower limb cellulitis. *J Am Acad Dermatol* 2012; **67**:163 e1–12; quiz 75–6.
- NHS Digital. Hospital Admitted Patient Care Activity 2018–19. 19 September 2019. Available at: <https://digital.nhs.uk/data-and-information/publications/statistical/hospital-admitted-patient-care-activity/2018-19>.
- Australian Institute of Health and Welfare. Potentially preventable hospitalisations in Australia by age groups and small geographic areas, 2017–18. 14 November 2019. Available at: <https://www.aihw.gov.au/reports/primary-health-care/potentially-preventable-hospitalisations/data>.
- Quere I, Palmier S, Noerregaard S *et al.* LIMPRINT: estimation of the prevalence of lymphoedema/chronic oedema in acute hospital in in-patients. *Lymphat Res Biol* 2019; **17**:135–40.
- Moffatt CJ, Gaskin R, Sykorova M *et al.* Prevalence and risk factors for chronic edema in U.K. community nursing services. *Lymphat Res Biol* 2019; **17**:147–54.
- Moffatt C, Keeley V, Quere I. The concept of chronic edema – a neglected public health issue and an international response: the LIMPRINT Study. *Lymphat Res Biol* 2019; **17**:121–6.
- Moffatt CJ, Franks PJ, Doherty DC *et al.* Lymphoedema: an underestimated health problem. *QJM* 2003; **96**:731–8.
- Quirke M, Ayoub F, McCabe A *et al.* Risk factors for nonpurulent leg cellulitis: a systematic review and meta-analysis. *Br J Dermatol* 2017; **177**:382–94.
- Al-Niaimi F, Cox N. Cellulitis and lymphoedema: a vicious cycle. *J Lymphoedema* 2009; **4**:38–42.
- Cox NH. Oedema as a risk factor for multiple episodes of cellulitis/erysipelas of the lower leg: a series with community follow-up. *Br J Dermatol* 2006; **155**:947–50.
- Dai M, Sugama J, Tsuchiya S *et al.* Inter-rater reliability of the AFTD-pitting test among elderly patients in a long-term medical facility. *Lymphoedema Res Pract* 2015; **3**:1–7.
- International Society of Lymphology. The diagnosis and treatment of peripheral lymphedema: 2013 Consensus document of the International Society of Lymphology. *Lymphology* 2013; **46**:1–11.
- Moffatt C, Franks P, Keeley V *et al.* The development and validation of the LIMPRINT methodology. *Lymphat Res Biol* 2019; **17**:127–34.
- Keeley V, Franks P, Quere I *et al.* LIMPRINT in specialist lymphedema services in United Kingdom, France, Italy, and Turkey. *Lymphat Res Biol* 2019; **17**:141–6.
- Moffatt CJ, Keeley V, Hughes A *et al.* LIMPRINT: the UK experience – subjective control of swelling in patients attending specialist lymphedema services. *Lymphat Res Biol* 2019; **17**:211–20.
- Dean SM, Valenti E, Hock K *et al.* The clinical characteristics of lower extremity lymphedema in 440 patients. *J Vasc Surg Venous Lymphat Disord* 2020; **8**:851–9.
- Rodriguez JR, Hsieh F, Huang CT *et al.* Clinical features, microbiological epidemiology and recommendations for management of cellulitis in extremity lymphedema. *J Surg Oncol* 2020; **121**:25–36.
- Park SI, Yang EJ, Kim DK *et al.* Prevalence and epidemiological factors involved in cellulitis in Korean patients with lymphedema. *Ann Rehabil Med* 2016; **40**:326–33.
- Teerachaisakul M, Ekataksin W, Durongwatana S *et al.* Risk factors for cellulitis in patients with lymphedema: a case-controlled study. *Lymphology* 2013; **46**:150–6.
- Lymphoedema Framework. Best Practice for the Management of Lymphoedema. International consensus. London: MEP Ltd, 2006. Available at: https://www.lympho.org/wp-content/uploads/2016/03/Best_practice.pdf.
- Dalal A, Eskin-Schwartz M, Mimouni D *et al.* Interventions for the prevention of recurrent erysipelas and cellulitis. *Cochrane Database System Rev* 2017; **6**:CD009758.

- 23 Webb E, Neeman T, Bowden FJ *et al.* Compression therapy to prevent recurrent cellulitis of the leg. *N Engl J Med* 2020; **383**:630–9.
- 24 Moffatt CJ, Doherty DC, Franks PJ *et al.* Community-based treatment for chronic edema: an effective service model. *Lymphat Res Biol* 2018; **16**:92–9.
- 25 Foldi E, Sauerwald A, Hennig B. Effect of complex decongestive physiotherapy on gene expression for the inflammatory response in peripheral lymphedema. *Lymphology* 2000; **33**:19–23.
- 26 International Lymphoedema Framework in association with the World Alliance for Wound and Lymphoedema Care. *Best Practice for the Management of Lymphoedema – 2nd Edition. Compression Therapy: A position document on compression bandaging*. 2012. Available at: <https://www.lympho.org/wp-content/uploads/2016/03/Compression-bandaging-final.pdf>.
- 27 Jones D, Meijer EFJ, Blatter C *et al.* Methicillin-resistant *Staphylococcus aureus* causes sustained collecting lymphatic vessel dysfunction. *Sci Transl Med* 2018; **10**:eaam7964.
- 28 Andersen CJ, Murphy KE, Fernandez ML. Impact of obesity and metabolic syndrome on immunity. *Adv Nutr* 2016; **7**:66–75.
- 29 Weitman ES, Aschen SZ, Farias-Eisner G *et al.* Obesity impairs lymphatic fluid transport and dendritic cell migration to lymph nodes. *PLOS ONE* 2013; **8**:e70703.
- 30 vom Steeg LG, Klein SL. Sex matters in infectious disease pathogenesis. *PLoS Pathog* 2016; **12**:e1005374.
- 31 Klein SL, Flanagan KL. Sex differences in immune responses. *Nat Rev Immunol* 2016; **16**:626–38.
- 32 Carey IM, Critchley JA, DeWilde S *et al.* Risk of infection in type 1 and type 2 diabetes compared with the general population: a matched cohort study. *Diabetes Care* 2018; **41**:513–21.
- 33 Ko LN, Garza-Mayers AC, St John J *et al.* Effect of dermatology consultation on outcomes for patients with presumed cellulitis: a randomized clinical trial. *JAMA Dermatol* 2018; **154**:529–36.

Appendix Conflicts of interest

All authors have completed the ICMJE uniform disclosure form at www.icmje.org/coi_disclosure.pdf and declare no support from any organization for the submitted work, except from P.J.F., who received funding from the International Lymphoedema Framework (ILF) during the conduct of the study. Reponex Pharmaceuticals and Coloplast sponsors E.A.B. in her PhD, through payments to the department, outside the submitted work. E.A.B. has also been an investigator for clinical research for Genentech, Reaplix, Ilkos Therapeutic and SoftOx Solutions, outside the submitted work. C.J.M. is sponsored by Thuasne and Essity Healthcare for consulting in compression therapy, and by the ILF for work on different research, outside the submitted work. T.K. is sponsored by Coloplast regarding stomas and wound healing as part of an advisory board membership, has been an investigator for clinical research for Genentech, and is a medical advisor for Reponex Pharmaceuticals, outside the submitted work. P.J.F. has received grants from Tactile Medical, outside the submitted work. I.Q. has received honoraria for consulting and as a speaker from Thuasne in the last 3 years. I.Q. was an investigator for clinical research for Thuasne and Medi; fees were paid to the hospital. V.K. was an investigator for clinical research for Tactile Medical outside the submitted work, with fees paid to the hospital. V.K. was a consultant to Koya and received an honorarium as a speaker from Medi. There are no other relationships or activities that could appear to have influenced the submitted work.