

Journal Pre-proof

Relationship Between Kinesiophobia and Quality of Life Among Patients with Breast Cancer-Related Lymphedema: Chain-Mediating Effect of Self-Care and Functional Exercise Compliance

Qi Wang, Na Du



PII: S2347-5625(23)00164-6

DOI: <https://doi.org/10.1016/j.apjon.2023.100346>

Reference: APJON 100346

To appear in: *Asia-Pacific Journal of Oncology Nursing*

Received Date: 1 September 2023

Revised Date: 14 November 2023

Accepted Date: 16 November 2023

Please cite this article as: Wang Q, Du N, Relationship Between Kinesiophobia and Quality of Life Among Patients with Breast Cancer-Related Lymphedema: Chain-Mediating Effect of Self-Care and Functional Exercise Compliance, *Asia-Pacific Journal of Oncology Nursing*, <https://doi.org/10.1016/j.apjon.2023.100346>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier Inc. on behalf of Asian Oncology Nursing Society.

Article type: Original article

The Relationship Between Kinesiophobia and Quality of Life Among Patients with Breast Cancer-Related Lymphedema: Chain-Mediating Effect of Self-Care and Functional Exercise Compliance

Qi Wang, Na Du*

Department of Gynaecology and Obstetrics, Shenging Hospital of China Medical University, 36 Sanhao St, Heping District, Shenyang 110000, Liaoning, China.

***Corresponding to:** Na Du, Department of Gynaecology and Obstetrics, Shenging Hospital of China Medical University, 36 Sanhao St, Heping District, Shenyang 110000, Liaoning, China.

Email address: Nanad_gny@163.com

ORCID ID: <https://orcid.org/0009-0001-0083-3323>

Statement and Declaration

Acknowledgement

The authors appreciate the research interviewers and the involved patients who generously gave of their own time to participate in this study.

CRedit author statement

Qi Wang: Methodology, Software, Data curation, Data analysis, Writing-Original draft preparation; **Na Du:** Research conceptualization, Validation, Supervision, Writing-Reviewing and Editing. All authors had full access to all the data in the study, and the corresponding author had final responsibility for the decision to submit for publication. The corresponding author attests that all listed authors meet authorship criteria and that

no others meeting the criteria have been omitted.

Ethics statement

This study was approved by the Research Ethics Committee of Shengjing Hospital Affiliated to China Medical University (No.EC-2020-HS-030), and in accordance with the Declaration of Helsinki. All participants provided written informed consent.

Funding

This study received no external funding.

Declaration of competing interest

The authors declare no conflict of interest.

Data availability statement

The data that support the findings of this study are available from the corresponding author ND, upon reasonable request.

Declaration of generative AI in scientific writing

No AI tools/services were used during the preparation of this manuscript.

1 **Relationship Between Kinesiophobia and Quality of Life Among Patients with Breast**
2 **Cancer-Related Lymphedema: Chain-Mediating Effect of Self-Care and Functional**
3 **Exercise Compliance**

4 **Abstract**

5 **Objective:** Breast cancer-related lymphedema (BCRL) significantly impacts the quality of life
6 (QoL) of breast cancer survivors following treatment. This study explores the association
7 between kinesiophobia (fear of pain caused by movement) and QoL in post-surgical BCRL
8 survivors and examines whether self-care and compliance with functional exercise act as
9 mediators between these variables.

10 **Methods:** This cross-sectional study surveyed 274 BCRL patients at three tertiary hospitals in
11 Shenyang City, China, from May 2020 to October 2022. The participants completed self-
12 reported questionnaires on self-care, functional exercise compliance, kinesiophobia, and QoL.
13 Medication analysis was conducted using the PROCESS Macro (Model 6).

14 **Results:** Kinesiophobia was found to have negative association with self-care ($p < 0.001$),
15 functional exercise compliance ($p < 0.001$), and QoL ($p < 0.001$). Kinesiophobia indirectly
16 affected QoL through three mediating pathways: self-care (effect = -0.132), functional exercise
17 compliance (effect = -0.390), and a combination of self-care and functional exercise
18 compliance (effect = -0.220), collectively accounting for 7.9%, 23.3%, and 13.1% of the total
19 effect, respectively.

20 **Conclusions:** This study highlights the substantial chain-mediating role of self-care and
21 functional exercise compliance in the relationship between kinesiophobia and QoL. It provides

- 22 valuable evidence supporting the protective effects of self-care and functional exercise
23 compliance in mitigating kinesiophobia and enhancing the QoL of BCRL survivors.
- 24 **Keywords:** Breast cancer-related lymphedema; Chain-mediated effect; Functional exercise
25 compliance; Kinesiophobia; Quality of life; Self-care

Journal Pre-proof

26 1. Introduction

27 Breast cancer (BC) is recognized as the most prevalent malignancy affecting women in
28 China.¹ The primary treatment for clinical BC often involves modified radical mastectomy
29 (MRM) combined with chemoradiotherapy. However, BC survivors frequently endure a
30 substantial burden of postoperative symptoms in the long term. Among these complications,
31 breast cancer-related lymphedema (BCRL) is one of the most common, affecting around 5%
32 to 50% of BC patients.² BC surgical procedures can disrupt the normal functioning of the
33 lymph system, leading to the accumulation of excess fluid in the affected upper extremity. This
34 condition is typically associated with severe swelling, pain, stiffness, and even limitations in
35 shoulder and arm mobility.³ Consequently, previous research has shown that the persistent
36 upper extremity impairments caused by BCRL not only severely impact patients' daily physical
37 activities, but also exacerbate their psychological distress, ultimately diminishing their overall
38 quality of life (QoL).⁴

39 QoL is a crucial index for assessing the multidimensional aspects of patients' health
40 outcomes in their daily lives during long-term rehabilitation of cancer survivors.⁵ Kwan et al.
41 verified that nearly 30% to 70% of BC patients experienced at least mild shoulder or arm
42 symptoms after BC treatment, and 12.5% of patients developed lymphedema, as revealed
43 through a survey of 744 BC patients. Furthermore, both symptomatic patients and BC patients
44 with lymphedema had significantly worse QoL scores compared to asymptomatic patients.⁶ In
45 addition, a cross-sectional study conducted by Bulley et al. found that the prevalence of
46 lymphedema peaked at 28% in the third year among 473 women who had undergone BC
47 treatment. Moreover, the study revealed that persistent arm problems seriously impaired

48 patients' upper limb function and led to a decline in their QoL over time.⁷ A recent review has
49 shown that patients with BCRL and poor QoL often experience feelings of weakness, financial
50 burden, concerns about disease progression, changes in body image, and reduced limb
51 function.⁸ Therefore, considering that the essential aspects of QoL such as physical functioning,
52 psychological health, social support, and spiritual well-being experienced by BC patients are
53 strongly associated with lymphedema, QoL is widely perceived as a particularly significant
54 factor in measuring symptom relief and prognosis among BC survivors.⁹

55 Kinesiophobia refers to a situation where individuals have an irrational and intense fear of
56 physical activity to avoid painful re-injury.¹⁰ Based on the fear avoidance model, this fear of
57 pain leads kinesiophobic patients to reduce their daily activities and physical movements,
58 ultimately resulting in the degradation of somatic functions and deterioration of mental
59 health.¹¹ Notably, Gencay et al. discovered that 76% of patients with kinesiophobia also had
60 upper extremity lymphedema among 81 BC patients who had undergone cancer-related surgery.
61 This suggests that those who develop lymphedema are more likely to limit the use and
62 movement of their upper extremities due to the fear of excessive pain, which can worsen the
63 severity of the edema.¹² Similarly, Karadibak et al. found that the severity of edema in upper
64 extremity lymphedema was positively correlated with a high level of kinesiophobia, but
65 negatively correlated with better QoL among 62 BC patients (all $p < 0.01$).¹³ Furthermore, a
66 cross-sectional study of 54 BC survivors reported that a high incidence (66.7%) of
67 kinesiophobia was significantly correlated with poorer QoL scores among these patients.¹⁴
68 Therefore, kinesiophobia is considered to play an essential role in predicting QoL in patients

69 with chronic disorders.¹⁵ However, the impact of kinesiophobia on QoL in BC patients with
70 lymphedema remains rarely studied.

71 Self-care may play a role in the interaction between kinesiophobia and QoL in BC patients
72 with lymphedema. Self-care ability is recognized as a series of conscious activities that can
73 assist cancer survivors in returning to their normal lives after surgery or chemoradiotherapy by
74 enhancing proactive attitudes and voluntarily modifying health-related behaviors.¹⁶ A prior
75 study revealed that low adherence to BCRL self-care modalities could accelerate the
76 development and progression of BCRL into more advanced stages. Therefore, it is imperative
77 to enhance the self-care abilities of these survivors to promote their QoL.¹⁷ Additionally,
78 according to the Health Belief Model, individuals with kinesiophobia tend to subjectively
79 amplify obstacles they may encounter; consequently, they restrict their health-related behaviors,
80 such as self-care.¹⁸ From this perspective, BC patients with postoperative upper extremity
81 lymphedema are advised to acquire adequate self-care practices from healthcare professionals
82 or nurses to enhance their management skills for BCRL and improve their QoL.¹⁹

83 Notably, previous researchers have suggested that BC survivors should actively engage in
84 postoperative functional exercise programs to prevent the progression of lymphedema in the
85 upper extremities.²⁰ However, it is a great challenge for the majority of patients to adhere to
86 long-term functional exercise programs after discharge.²¹ Comparable studies have illustrated
87 that the adherence rate to functional exercise gradually decreases after discharge, especially
88 from the 105th day post-operation, which directly hinders the recovery of normal upper limb
89 function.²² Accordingly, increasing numbers of scholars have demonstrated that fear of pain
90 and movement is the major barrier for BC survivors to utilize rehabilitation exercise.²³

91 Interestingly, a randomized controlled trial study revealed that BC patients who underwent 8
92 weeks of rehabilitative exercise intervention following surgery showed a significant
93 improvement in kinesiophobia scores compared to those in the control group.²⁴ Moreover,
94 Keradibak et al. reported that adherence to a 12-week home-based exercise program had
95 significantly positive effects on improving kinesiophobia and QoL in patients with BCRL
96 following BC surgery.¹³ However, the correlation between kinesiophobia, postoperative
97 exercise adherence, and QoL among survivors with BCRL has not yet been fully determined.

98 Self-care and adherence to functional exercises are closely interconnected, especially in
99 cases of chronic illnesses. Warehime et al. conducted a qualitative study that substantiated the
100 importance of confidently participating in self-care activities in patients with heart failure. The
101 study found that 59.1% of patients who engaged in self-care activities experienced improved
102 health outcomes and long-term exercise adherence, suggesting that self-care might play a pivotal
103 role in adherence to functional exercise programs.²⁵ Interestingly, a randomized controlled
104 study confirmed that BC patients in the intervention group who participated in a 10-minute
105 holistic BCRL self-care program including gentle arm functional exercises for 6 months,
106 exhibited higher improvements in BCRL-related symptoms, self-care scores, and exercise
107 frequency (all $p < 0.05$) compared to those in the control group, highlighting the beneficial
108 effects of the BCRL self-care program among BC patients.²⁶ Additionally, a small sample
109 feasibility study confirmed that patients with BCRL who followed daily home-based exercise
110 combining with standard lymphedema self-care measures for 26 weeks experienced a clinically
111 meaningful improvement in the management of BCRL, adherence to the exercise program, and
112 QoL, compared to the control group that only practiced self-care measures.²⁷

113 Currently, there has been no evidence regarding the evaluation of the underlying
114 mechanisms of self-care ability, compliance with functional exercise, kinesiophobia, or QoL in
115 patients with BCRL. Therefore, this study aimed to determine the correlation between
116 kinesiophobia and QoL, as well as the underlying mechanical roles of self-care ability and
117 functional exercise compliance among BC patients with lymphedema. We hypothesized that
118 (1) kinesiophobia would negatively predict and directly affect QoL; (2) self-care would act as
119 a mediator in the relationship between kinesiophobia and QoL; (3) functional exercise
120 compliance would play an intermediary role between kinesiophobia and QoL; and (4) both
121 self-care and functional exercise compliance would serve as chain mediators in the correlation
122 between kinesiophobia and QoL. The hypothesized framework is shown in Fig. 1 and provides
123 valuable evidence for the establishment of targeted interventions to improve the QoL of BC
124 survivors with lymphedema.

125 **2. Methods**

126 **2.1. Participants and design**

127 A cross-sectional study was conducted by surveying 300 patients with BCRL from three
128 tertiary hospitals in Shenyang City, China, between May 2020 and October 2022. The inclusion
129 criteria were as follows: (1) female patients diagnosed with BCRL; (2) at least 60 days after
130 completing their cancer-related treatments (including surgical procedures and postoperative
131 chemotherapy or radiotherapy); (3) had clear cognition and normal communication ability; (4)
132 aged > 18 years old; and (5) no preexisting upper extremity dysfunction before surgery. The
133 exclusion criteria were as follows: (1) psychiatric disorders and poor cooperation; (2) other

134 malignancies or breast diseases; (3) liver, heart, or kidney failure; and (4) had undergone
135 bilateral axillary lymphadenectomy.

136 **2.2. Data collection and procedure**

137 This study was conducted by qualified researchers who had received professional training
138 at each outpatient oncology clinic in the three tertiary hospitals. The participants were enrolled
139 using convenience sampling. To avoid potential response bias, all participants were
140 individually invited to participate in the survey in a private and quiet area within the nurses'
141 station without the companionship of family members or friends. At the outset, the purpose and
142 methodology of this investigation were thoroughly explained to each patient, and the
143 participants were assured that they could withdraw from the survey at any time without any
144 impact on their treatment. The paper-based Chinese version of the questionnaires were
145 distributed to participants by the researchers. Each respondent completed the questionnaire
146 both anonymously and independently. During the investigation, the researchers were alongside
147 the participants to help them understand any confusing survey items. It took the participants
148 approximately 10 to 15 minutes to complete all the self-reported questionnaires. After the
149 survey, trained researchers immediately checked each participant's responses to guarantee that
150 all the required data were filled out. All the participants were informed that the collected data
151 would be confidential and would only be used for research purposes.

152 **2.3. Sample size**

153 The sample size was estimated by applying a metric of 5 to 10 respondents per item in a
154 validated survey to ensure sufficient statistical power.^{28, 29} The instrument used in this study
155 with the largest number of items was the Chinese Functional Assessment of Cancer Therapy-

156 Breast version 4.0 (FACT-Bv4.0), which consists of 36 items. The required sample size was
157 180 participants. Considering a sample loss of 20%, a final sample size of 225 participants was
158 required. We approached 315 patients with BCRL, of whom 300 patients met the inclusion
159 criteria and agreed to participate in the study (response rate, 95.2%). Among the 300 eligible
160 participants, 26 were excluded because of withdrawal (7 cases) or missing data (19 cases).
161 Ultimately, 274 completed questionnaires were collected for final analysis, with a valid
162 recovery rate of 91.3% (Fig. 2).

163 **2.4. Measurements**

164 **2.4.1. Demographic and clinical characteristics**

165 The demographic data of the participants, including age, body mass index (BMI),
166 educational level, marital and employment status, monthly family income, and number of
167 children, and clinical data, such as disease duration, BC stage, treatment type, tumor location,
168 and comorbidities (e.g., hypertension, diabetes, thyroid, osteoporosis, or dyslipidemia), were
169 obtained either from self-designed surveys or available medical records in the hospitals by the
170 responsible nursing staff.

171 **2.4.2. Measurement of self-care ability**

172 Self-care ability was evaluated by the Appraisal of Self-Care Agency Scale-Revised
173 (ASAS-R), which contains 15 items and is classified into three dimensions: having self-care
174 capacity, developing self-care capacity, and lack of self-care capacity, of which the lack of self-
175 care capacity was adversely recorded. Total scores range from 15 to 75 using a 5-point Likert
176 type from 1 (totally disagree) to 5 (totally agree). Higher scores indicate better self-care. The
177 Cronbach's α coefficient in this study was 0.80.

178 **2.4.3. Evaluation of functional exercise adherence**

179 The Postoperative Functional Exercise Adherence Scale (PFEAS) was used to measure
180 functional exercise adherence according to previous Chinese researchers.³⁰ This self-report
181 scale consists of 18 items on a 4-point Likert scale, ranging from 1 (unable to accomplish) to
182 4 (competent to accomplish). The scale is divided into three domains: postoperative
183 precautionary adherence, physical exercise adherence, and actively seeking advice adherence.
184 The total score ranges from 18 to 72, with higher scores indicating higher levels of adherence.
185 The Cronbach's α coefficient in this study was 0.87.

186 **2.4.4. Measurement of Kinesiophobia**

187 Kinesiophobia was measured using the Tampa Scale for Kinesiophobia-11 (TSK-11). It
188 comprises 11 items rated on a 4-point Likert-type scale, ranging from 1 (completely disagree)
189 to 4 (completely agree), to assess the patients' fear of movement or reinjury related to pain.
190 The total score ranged from 11 to 44, with a higher score indicating a higher level of
191 kinesiophobia: a score of ≤ 17 indicated no fear of movement; a score of 18-24 indicated mild
192 fear; a score of 25-31 indicated moderate fear; a score of 32-38 indicated severe fear, and a
193 score of 39-44 indicated extreme fear. The Cronbach's α coefficient in this study was 0.82.

194 **2.4.5. Assessment of QoL**

195 FACT-Bv4.0 was used to evaluate the QoL, which consists of 36 items rated on a 5-point
196 Likert scale from 0 (totally disagree) to 4 (totally agree), with total scores ranging from 0 to
197 144. The assessment comprises five subscales: a general cancer subscale (FACT-G), including
198 physical well-being (PWB, seven items), emotional well-being (EWB, six items), social/family
199 well-being (SWB, seven items), and functional well-being (FWB, seven items); and a Breast

200 Cancer Subscale for additional concerns (BCS, nine items). Higher scores indicate better QoL.
201 In this study, the Cronbach's α for the above subscales was 0.82 (PWB), 0.83 (EWB), 0.87
202 (SWB), 0.80 (FWB), and 0.87 (BCS), and the Cronbach's α for the overall scale was 0.91.

203 **2.5. Data analysis**

204 All data were analyzed using SPSS Statistics software version 25 (IBM, Inc., Chicago, IL,
205 USA). The normal distribution of the variables was tested using the Kolmogorov-Smirnov
206 method. Continuous variables were expressed as mean \pm standard deviation (SD), while
207 categorical data were presented as frequencies with percentages (%). Independent *t*-test and
208 one-way analysis of variance (ANOVA) were performed to determine the statistical
209 significance of the different demographic and clinical groups. Harman's single-factor test was
210 employed to mitigate the possibility of common method bias underlying the observed results,
211 which could be attributed to the single source of data collection in our study.³¹ Pearson's
212 correlation analysis was performed to examine the relationships between kinesiophobia, self-
213 care, functional exercise compliance, and QoL. Model 6 of the PROCESS Macro was
214 employed to estimate the chain-mediating effect of self-care and functional exercise adherence
215 (two mediators) in the relationship between kinesiophobia (one independent variable) and QoL
216 (one dependent variable). Demographic and clinical characteristics, including age, marital
217 status, BC duration, BC stage, chemotherapy, radiotherapy, and comorbidities that had a
218 significant impact on kinesiophobia, self-care, functional exercise adherence, and QoL (Table
219 1) were incorporated into the model as control variables. Bootstrapping was performed using
220 5000 random samples to determine the significance of the mediating effect. If the
221 corresponding 95% bias-corrected confidence interval (CI) did not contain zero, the mediating

222 effect was considered statistically significant. A value of $p < 0.05$ (two-tailed) was considered
223 statistically significant.

224 **2.6. Ethical considerations**

225 This study was approved by the Human Research Ethical Committee of Shengjing
226 Hospital of China Medical University (No.EC-2020-HS-030) and was conducted in accordance
227 with the Declaration of Helsinki. This study adhered to the Strengthening the Reporting of
228 Observation Studies in Epidemiology (STROBE). Written informed consent was obtained from
229 each respondent before they participated in the study.

230 **3. Results**

231 **3.1. Common method bias test**

232 Harman's single-factor test was used to detect possible common methodological biases.
233 The results showed that there were 22 factors with characteristic values greater than 1, and the
234 amount of variance explained by the first factor was 21.6%, which was less than the critical
235 criterion of 40%. Thus, the influence of common methodological deviations was excluded from
236 this study.

237 **3.2. Different variable scores according to characteristics of participants**

238 As shown in Table 1, the mean age of the participants was 48.76 ± 8.78 years old, ranging
239 from 27 to 76 years. More than half of the patients (55.1%) had been diagnosed with BC for
240 more than 4 years. Most patients were married (84.3%) and had children (86.9%). Nearly 44.2%
241 were employed and 35.4% had obtained a high school education degree or higher. The majority
242 of patients (77.4%) had T1 and TII stages, and approximately 62.4% of patients underwent
243 MRM therapeutic operation, while 30.7% underwent breast-protective surgery (BPS).

244 Approximately 55.8% of the patients received chemotherapy, and 50.4% underwent
245 radiotherapy. The other demographic and clinical characteristics were presented in Table 1.
246 Moreover, age ($p = 0.038$), marital status ($p = 0.007$), BC duration ($p = 0.015$), BC stage ($p =$
247 0.003), chemotherapy ($p = 0.008$), radiotherapy ($p = 0.006$), and comorbidities ($p = 0.040$)
248 were significantly associated with kinesiophobia (Table 1). Furthermore, older adult patients
249 ($p = 0.002$), those with prolonged BC duration ($p = 0.004$), and those who underwent
250 radiotherapy ($p = 0.019$) exhibited lower self-care scores than those who did not (Table 1).
251 Additionally, older adult patients ($p = 0.001$) who underwent chemotherapy ($p = 0.013$) were
252 less likely to adhere to postoperative functional exercises (Table 1). Regarding QoL, patients
253 who were older ($p = 0.004$), single ($p = 0.035$), had a prolonged BC duration ($p = 0.021$), had
254 an advanced BC stage ($p = 0.010$), received chemotherapy ($p = 0.006$) or radiotherapy ($p <$
255 0.001), and had comorbidities ($p = 0.042$) displayed worse QoL among these participants
256 (Table 1).

257 **3.3. Correlation analyses among the investigated variables**

258 As shown in Table 2, the total scores of kinesiophobia, self-care, functional exercise
259 adherence, and QoL were 29.27 ± 4.49 , 40.95 ± 5.03 , 43.09 ± 7.29 , and 79.52 ± 11.93 ,
260 respectively (Table 2). Pearson's correlation analysis showed that kinesiophobia was
261 negatively associated with self-care ($r = -0.481$, $p < 0.001$), functional exercise adherence ($r =$
262 -0.564 , $p < 0.001$), and QoL ($r = -0.680$, $p < 0.001$; Table 2). Additionally, self-care exhibited
263 a significantly positive correlation with functional exercise adherence ($r = 0.614$, $p < 0.001$)
264 and QoL ($r = 0.569$, $p < 0.001$; Table 2). A strong positive correlation was observed between
265 adherence to functional exercise and QoL ($r = 0.713$, $p < 0.001$; Table 2).

266 **3.4. Mediating effect of self-care and functional exercise adherence**

267 Considering QoL as the dependent variable, kinesiophobia as the independent variable, and
268 self-care and functional exercise adherence as the intermediary variables, the chain mediation
269 model of the mediating effects of self-care and functional exercise adherence between
270 kinesiophobia and QoL was shown in Fig. 2 and Table 3. Demographic and clinical
271 characteristics such as age, marital status, BC duration, BC stage, chemotherapy, radiotherapy,
272 and comorbidities were included as control variables. The regression analysis showed that the
273 total effect of kinesiophobia on QoL was significant ($\beta = -0.630$, $t = -13.276$, $p < 0.001$; Fig. 2
274 and Table 3). Furthermore, there was a significantly direct effect of kinesiophobia on QoL (β
275 $= -0.351$, $t = -7.407$, $p < 0.001$; Fig. 2 and Table 3), indicating that the higher levels of
276 kinesiophobia were associated with worse QoL among BC patients with lymphedema.
277 Moreover, kinesiophobia had a significant negative predictive effect on self-care ($\beta = -0.441$, t
278 $= -7.751$, $p < 0.001$), and self-care positively predicted QoL ($\beta = 0.113$, $t = 2.322$, $p < 0.05$; Fig.
279 2 and Table 3), suggesting the mediating role of self-care played between kinesiophobia and
280 QoL. Meanwhile, kinesiophobia had a negative impact on functional exercise adherence ($\beta =$
281 -0.349 , $t = -6.598$, $p < 0.001$), while functional exercise adherence had a positive effect on QoL
282 ($\beta = 0.420$, $t = 8.237$, $p < 0.001$; Fig. 2 and Table 3), implying that functional exercise adherence
283 acted as a mediator between kinesiophobia and QoL. Furthermore, self-care exhibited a
284 positive correlation with functional exercise adherence ($\beta = 0.446$, $t = 8.650$, $p < 0.001$; Fig. 2
285 and Table 3), indicating that self-care and functional exercise adherence played a chain
286 mediating role between kinesiophobia and QoL of BC survivors with lymphedema.

287 **3.5. Bootstrap examination**

288 We validated the mediating effect using a Bootstrap approach. The 95% CI for the
289 mediating effect was calculated based on a randomly selected sample of 5000 individuals to
290 assess the mediating role of self-care and functional exercise adherence as mediators between
291 kinesiophobia and QoL. As shown in Table 4, the direct effect pathway was as follows:
292 kinesiophobia \rightarrow QoL, with a direct effect value of -0.933 (Bootstrap 95% CI: -1.181 to -0.685),
293 which accounted for 55.7% of the total effect (Table 4). The total indirect effect was -0.742
294 (Bootstrap 95% CI: -0.915 to -0.592), accounting for 44.3% of the total effect (Table 4). The
295 lower and upper Bootstrap 95% CI did not contain a value of zero, indicating that the effect
296 was significant. Meanwhile, there were three mediating effect pathways: kinesiophobia \rightarrow self-
297 care \rightarrow QoL, with an indirect effect value of -0.132 (Bootstrap 95% CI: -0.267 to -0.007),
298 accounting for 7.9% of the total effect (Table 4); kinesiophobia \rightarrow functional exercise
299 adherence \rightarrow QoL, with an indirect effect value of 0.390 (Bootstrap 95% CI: -0.516 to -0.269),
300 accounting for 23.3% of the total effect (Table 4); and kinesiophobia \rightarrow self-care \rightarrow functional
301 exercise adherence \rightarrow QoL, with an indirect effect value of -0.220 (Bootstrap 95% CI: -0.320
302 to -0.142), accounting for 13.1% of the total effect (Table 4). Hence, Hypotheses 1, 2, 3, and 4
303 are verified.

304 **4. Discussion**

305 The majority of BC survivors tend to avoid using their affected arms after treatment
306 because of kinesiophobia, which results in an increased risk of developing BCRL in the upper
307 extremity.¹² An accumulating stream of studies has demonstrated that sufficient self-care
308 capacity and adherence to functional exercises can help improve edema and have a potent effect
309 on the QoL of women with lymphedema.^{26, 27} Nevertheless, the relationship between

310 kinesiophobia and QoL has rarely been examined in detail. Therefore, the present study aimed
311 to explore this hypothesized association, which may be connected to self-care and adherence
312 to functional exercise.

313 **4.1. Relationship between kinesiophobia and QoL**

314 Kinesiophobia is rather frequent and strongly correlated with lymphedema in BC survivors
315 after mastectomy.³² In the current study, we found that all 274 patients with BCRL included in
316 this study exhibited varying degrees of kinesiophobia (TSK-11 score \geq 18 points). Similarly,
317 Altas and Demirdel found that among 70 female patients with post-mastectomy lymphedema,
318 kinesiophobia was presented in 70% of these patients, as measured using the TSK.³² These
319 findings highlights the high prevalence of kinesiophobia among patients with BCRL after
320 undergoing BC surgery.¹² Furthermore, our study confirmed a correlation between fear of
321 movement and impaired QoL in patients with BCRL. We also discovered that kinesiophobia
322 served as a direct predictor of QoL in BC survivors with lymphedema, as higher kinesiophobia
323 scores were negatively associated with better QoL among patients with BCRL (Hypothesis 1
324 was confirmed). These findings were consistent with those of several previous studies. For
325 instance, a cross-sectional study conducted by Sunar found a significant correlation between a
326 high score of kinesiophobia and worse QoL, and fatigue was found among BC patients from
327 Turkey.¹⁴ Moreover, Gencay et al. confirmed that kinesiophobic patients had a significantly
328 lower physical QoL score in BC patients, although no significant correlation between
329 kinesiophobia and QoL scores was found.¹² Therefore, it is imperative for patients with BCRL
330 to receive appropriate training to help them promote pain management and overcome the fear
331 of movement, as well as improve their QoL. For example, a longitudinal cohort study

332 conducted by Velthuis et al. found that a high level of kinesiophobia was negatively associated
333 with the perceived global health status (QoL). However, this negative association was
334 significantly reduced by a 12-week graded activity rehabilitation program among cancer
335 survivors.³³ Therefore, professional medical staff should provide more practical and effective
336 support to help survivors overcome kinesiophobia and enhance their QoL.

337 **4.2. Mediating role of self-care on kinesiophobia and QoL**

338 The present study confirms Hypothesis 2, which states that self-care partially mediates the
339 relationship between kinesiophobia and QoL. The self-care ability of the patients with BCRL
340 in our study was relatively low, with a total score of 40.95 ± 5.03 . These findings were also
341 confirmed by Jiang et al., who developed factor-based models to demonstrate that the severity
342 of lymphedema was associated with decreased limb activity and lower self-care ability in
343 patients with BCRL.³⁴ Therefore, it is necessary for patients with BCRL to adhere to
344 lymphedema self-care behaviors to improve lymph drainage capacity and enhance their QoL.
345 Unfortunately, the overall self-care capacity of patients at risk of lymphedema is considerably
346 poor, and the reasons for non-adherence to self-care consist of symptom burden, complicated
347 treatment regimens, and insufficient educational support.³⁵ Therefore, Tsuchiya et al.
348 suggested that in addition to providing basic instructions on self-care skills, it was essential to
349 implement continuous psycho-educational programs to encourage female cancer survivors to
350 engage in self-care behaviors after discharge.³⁶ Specifically, to improve the QoL of BC
351 survivors with lymphedema, enhancing patients' awareness by educating them about the risk
352 of lymphedema and knowledge about proper self-care behaviors for BC cancer survivors is
353 warranted.³⁷

354 **4.3. Mediating role of functional exercise adherence on kinesiophobia and QoL**

355 The mediation model also showed that functional exercise adherence exhibited a significant
356 indirect effect on the relationship between kinesiophobia and QoL (Hypothesis 3 was verified).
357 After reviewing several papers on health-related QoL in BC published in the last decade,
358 scholars have summarized that appropriate physical activity interventions are effective in
359 managing symptoms of BCRL and improving QoL in BC survivors.³⁸ Nevertheless, previous
360 studies have suggested that women with BC often hold misconceptions that physical activity
361 can accelerate the spread of tumor cells and aggravate cancer metastases. Consequently, most
362 BC survivors refuse to adhere to the physical exercise recommendations provided by medical
363 professionals after treatment, which exacerbates the severity of BCRL.³⁹ In addition, it is
364 essential to provide proper education and implement a scientifically designed exercise program
365 to assist patients with BCRL in overcoming their fear of movement during rehabilitation. A
366 systematic review conducted by Baumann found that various types of exercise programs, such
367 as aerobic exercise, resistance exercise, aqua lymph training, yoga, and gravity-resistive
368 exercise, demonstrated a marked improvement in the severity of BCRL status, mood, and
369 QoL.²⁰ Cormie et al. conducted a randomized trial study in Australia to evaluate the safety and
370 efficacy of a prescribed resistance exercise program on 62 women with BCRL. They
371 discovered that patients who received appropriate upper-body resistance exercise intervention
372 for three months demonstrated significantly improved lymphedema management and higher
373 scores in the physical functioning domain of QoL than those in the usual care group (all $p <$
374 0.05).⁴⁰

375 **4.4. Chain-mediated role of self-care and functional exercise adherence on kinesiophobia**
376 **and QoL**

377 Regarding the relationship between kinesiophobia and QoL in patients with BCRL, this
378 study is the first to find that self-care and adherence to functional exercise have a chain-
379 mediating effect on the interaction between kinesiophobia and QoL in patients with BCRL.
380 The indirect effect ratio was 13.13%, supporting Hypothesis 4. These findings imply that the
381 fear of movement caused by a high level of kinesiophobia could impair self-care behaviors
382 among patients with BCRL. Their limited self-care abilities further prevent them from actively
383 participating in rehabilitative regimens, thereby resulting in low adherence to functional
384 exercise and decreased QoL. Conversely, patients with BCRL would benefit from adopting
385 adequate self-care behaviors, as this could assist them in overcoming the psychological
386 obstacles associated with kinesiophobia. Proper self-care behaviors also promote subjective
387 initiatives and perceived awareness of functional exercise adherence, thereby positively
388 influencing all dimensions of QoL among patients with BCRL. Recent randomized controlled
389 studies have emphasized the effectiveness of implementing proper self-care programs to
390 alleviate symptoms related to BCRL, promote exercise adherence, and improve QoL in patients
391 with BCRL.^{26, 27} Given the hypothesized framework used in our study, we found a positive
392 association between self-care and functional exercise adherence, suggesting that individuals
393 with higher self-care abilities are more likely to comply with functional exercise. This finding
394 was consistent with prior clinical studies conducted by Li et al.⁴¹ who demonstrated that the
395 implementation of self-controlled exercise programs could sustainably enhance the mobility of
396 the affected shoulder joints and have a positive effect on the post-operative recovery process

397 in BC patients. Furthermore, our findings confirmed the positive correlation between
398 functional exercise adherence and QoL, a relationship supported by other researchers. Kim et
399 al. found that participation in physical activity for self-care showed a significant positive
400 correlation with increased mobility, reduced mental burden, and improved QoL among Korean
401 patients with BC.⁴²

402 Notably, the chain mediating pathway of “self-care → functional exercise adherence” acted
403 as an essential bridge between kinesiophobia and QoL. Therefore, the results of the present
404 study extend our understanding of the potential variables that impact kinesiophobia and
405 uncover the pathways through which QoL can be promoted. These findings provide valuable
406 information for the development of effective clinical intervention strategies. Understanding the
407 psychological mechanisms in patients with BCRL contributes to promoting QoL by
408 emphasizing high-quality self-care practices and interventions that enhance adherence to
409 functional exercise. For instance, when addressing the challenges of kinesiophobia, nursing
410 staff should educate patients with BCRL on appropriate self-care practices that can improve
411 their health outcomes. Healthcare practitioners should provide tailored functional exercise
412 interventions during rehabilitation to enhance patient adherence. Moreover, effective
413 interventions that target the reduction of kinesiophobia can also improve patients’ self-care
414 abilities and enhance their motivation to engage in rehabilitative programs, ultimately resulting
415 in improved QoL among patients with BCRL. Additionally, when considering the chain-
416 mediating effects of self-care and functional exercise adherence, it is critical for oncology
417 nursing staff to develop comprehensive interventions that address this pathway: diminish
418 kinesiophobia → enhance self-care ability → promote functional exercise adherence →

419 improve QoL, which would be more effective in improving the health benefits of patients with
420 BCRL than interventions that target only one factor.

421 **4.5. Limitations**

422 This study has some limitations. First, the data were collected from only three urban
423 hospitals in the city, which may challenge the generalizability of the results. Extended surveys
424 are required to investigate patients with BCRL in rural regions and other cities in China. Second,
425 this study was conducted during the COVID-19 pandemic, which might have affected the
426 patients' self-estimated QoL and generated a response bias. Third, this study was cross-
427 sectional using self-reported data; thus, a longitudinal study should be conducted to further
428 identify the causal relationships of this framework and track changes in patients' attitudes
429 towards the aforementioned variables over the course of long-term cancer treatment. Fourth,
430 this study focused only on exploring the mechanism underlying the chain-mediating effects of
431 self-care capacity and functional exercise adherence on the relationship between kinesiophobia
432 and QoL in patients with BCRL. There may be other factors that affect QoL through different
433 mechanisms. Thus, a more comprehensive study should be conducted to further explore the
434 impact of kinesiophobia on the QoL of patients with BCRL.

435 **5. Conclusion**

436 To the best of our knowledge, the current study is the first to elucidate that kinesiophobia
437 has a negative impact on QoL in survivors of BCRL and the sequential mediating effect of self-
438 care and functional exercise adherence on the relationship between kinesiophobia and QoL in
439 patients with BCRL. The chain-mediating effect of self-care and adherence to functional
440 exercise represents potential practical significance in promoting the QoL of survivors with

441 BCRL. Therefore, clinical intervention programs that focus on improving self-care ability
442 should be developed and implemented to help diminish the detrimental impact of kinesiophobia
443 on QoL and expand the protective effects of adhering to functional exercise for survivors with
444 BCRL in the future.

Journal Pre-proof

445 **Statement and Declaration**

446 **Acknowledgement**

447 The authors appreciate the research interviewers and the involved patients who generously
448 gave of their own time to participate in this study.

449 **CRedit author statement**

450 **Qi Wang:** Methodology, Software, Data curation, Data analysis, Writing-Original draft
451 preparation; **Na Du:** Research conceptualization, Validation, Supervision, Writing-Reviewing
452 and Editing. All authors had full access to all the data in the study, and the corresponding author
453 had final responsibility for the decision to submit for publication. The corresponding author
454 attests that all listed authors meet authorship criteria and that no others meeting the criteria
455 have been omitted.

456 **Ethics statement**

457 This study was approved by the Research Ethics Committee of Shengjing Hospital Affiliated
458 to China Medical University (No.EC-2020-HS-030), and in accordance with the Declaration
459 of Helsinki. All participants provided written informed consent.

460 **Funding**

461 This study received no external funding.

462 **Declaration of competing interest**

463 The authors declare no conflict of interest.

464 **Data availability statement**

465 The data that support the findings of this study are available from the corresponding author
466 ND, upon reasonable request.

467 **Declaration of generative AI in scientific writing**

468 No AI tools/services were used during the preparation of this manuscript.

Journal Pre-proof

469 **References**

- 470 1. Wang X, Wang C, Guan J, Chen B, Xu L, Chen C. Progress of Breast Cancer basic research
471 in China. *Int J Biol Sci.* 2021;17:2069-2079. <https://doi.org/10.7150/ijbs.60631>.
- 472 2. Gillespie TC, Sayegh HE, Brunelle CL, Daniell KM, Taghian AG. Breast cancer-related
473 lymphedema: risk factors, precautionary measures, and treatments. *Gland Surg.* 2018;7:379-
474 403. <https://doi.org/10.21037/gs.2017.11.04>.
- 475 3. Shah C, Vicini FA. Breast cancer-related arm lymphedema: incidence rates, diagnostic
476 techniques, optimal management and risk reduction strategies. *Int J Radiat Oncol Biol Phys.*
477 2011;81:907-914. <https://doi.org/10.1016/j.ijrobp.2011.05.043>.
- 478 4. Nesvold IL, Fosså SD, Holm I, Naume B, Dahl AA. Arm/shoulder problems in breast cancer
479 survivors are associated with reduced health and poorer physical quality of life. *Acta Oncol.*
480 2010;49:347-353. <https://doi.org/10.3109/02841860903302905>.
- 481 5. Licu M, Ionescu CG, Paun S. Quality of Life in Cancer Patients: The Modern Psycho-
482 Oncologic Approach for Romania-A Review. *Curr Oncol.* 2023;30:6964-6975.
483 <https://doi.org/10.3390/currenol30070504>.
- 484 6. Kwan W, Jackson J, Weir LM, Dingee C, McGregor G, Olivotto IA. Chronic arm morbidity
485 after curative breast cancer treatment: prevalence and impact on quality of life. *J Clin Oncol.*
486 2002;20:4242-4248. <https://doi.org/10.1200/JCO.2002.09.018>.
- 487 7. Bulley C, Coutts F, Blyth C, et al. Prevalence and impacts of upper limb morbidity after
488 treatment for breast cancer: a cross-sectional study of lymphedema and function. *Cancer and*
489 *Oncology Research.* 2013;1:30-39. <https://doi.org/10.13189/cor.2013.010203>.
- 490 8. Kalemikerakis I, Evaggelakou A, Kavga A, Vastardi M, Konstantinidis T, Govina O.

- 491 Diagnosis, treatment and quality of life in patients with cancer-related lymphedema. *J BUON*.
492 2021;26:1735-1741. PMID: 34761576.
- 493 9. Abdo J, Ortman H, Rodriguez N, Tillman R, Riordan EO, Seydel A. Quality of Life Issues
494 Following Breast Cancer Treatment. *Surg Clin North Am*. 2023;103:155-167.
495 <https://doi.org/10.1016/j.suc.2022.08.014>.
- 496 10. Bordeleau M, Vincenot M, Lefevre S, et al. Treatments for kinesiophobia in people with
497 chronic pain: A scoping review. *Front Behav Neurosci*. 2022;16:933483.
498 <https://doi.org/10.3389/fnbeh.2022.933483>.
- 499 11. Vlaeyen JW, Kole-Snijders AM, Rotteveel AM, Ruesink R, Heuts PH. The role of fear of
500 movement/(re)injury in pain disability. *J Occup Rehabil*. 1995;5:235-252.
501 <https://doi.org/10.1007/BF02109988>.
- 502 12. Gencay Can A, Can SS, Ekşioğlu E, Çakıcı FA. Is kinesiophobia associated with
503 lymphedema, upper extremity function, and psychological morbidity in breast cancer survivors?
504 *Turk J Phys Med Rehabil*. 2018;65:139-146. <https://doi.org/10.5606/tftrd.2019.2585>.
- 505 13. Karadibak D, Yavuzsen T, Saydam S. Prospective trial of intensive decongestive
506 physiotherapy for upper extremity lymphedema. *J Surg Oncol*. 2008;97:572-577.
507 <https://doi.org/10.1002/jso.21035>.
- 508 14. Sunar İ, Sunar V. Kinesiophobia in Breast Cancer Survivors and its Relationship with
509 Quality of Life, Comorbidity, and other clinical parameters. *Acta Oncologica Turcica*.
510 2021;54:198-205. <https://doi.org/10.5505/aot.2021.38243>.
- 511 15. Nederhand MJ, Ijzerman MJ, Hermens HJ, Turk DC, Zilvold G. Predictive value of fear
512 avoidance in developing chronic neck pain disability: consequences for clinical decision

- 513 making. Arch Phys Med Rehabil. 2004;85:496-501.
514 <https://doi.org/10.1016/j.apmr.2003.06.019>.
- 515 16. Zhang Y, Kwekkeboom K, Petrini M. Uncertainty, Self-efficacy, and Self-care Behavior in
516 Patients With Breast Cancer Undergoing Chemotherapy in China. *Cancer Nurs*. 2015;38:E19-
517 26. <https://doi.org/10.1097/NCC.0000000000000165>.
- 518 17. Brown JC, Cheville AI, Tchou JC, Harris SR, Schmitz KH. Prescription and adherence to
519 lymphedema self-care modalities among women with breast cancer-related lymphedema.
520 *Support Care Cancer*. 2014;22:135-143. <https://doi.org/10.1007/s00520-013-1962-9>.
- 521 18. Zhang S, Wang Z, Lin X, et al. Kinesiophobia and self-management behaviour related to
522 physical activity in Chinese patients with coronary heart disease: The mediating role of self-
523 efficacy. *Nurs Open*. 2023;10:105-114. <https://doi.org/10.1002/nop2.1283>.
- 524 19. Deveci Z, Karayurt Ö, Eyigör S. Self-care practices, patient education in women with breast
525 cancer-related lymphedema. *Turk J Phys Med Rehabil*. 2021;67:187-195.
526 <https://doi.org/10.5606/tftrd.2021.5022>.
- 527 20. Baumann FT, Reike A, Reimer V, et al. Effects of physical exercise on breast cancer-related
528 secondary lymphedema: a systematic review. *Breast Cancer Res Treat*. 2018;170:1-13.
529 <https://doi.org/10.1007/s10549-018-4725-y>.
- 530 21. Tao L, Wang M, Zhang X, Du X, Fu L. Exercise adherence in breast cancer patients: A
531 cross-sectional questionnaire survey. *Medicine (Baltimore)*. 2020;99:e20427.
532 <https://doi.org/10.1097/MD.00000000000020427>.
- 533 22. Petito EL, Nazário AC, Martinelli SE, Facina G, De Gutiérrez MG. Application of a
534 domicile-based exercise program for shoulder rehabilitation after breast cancer surgery. *Rev*

- 535 Lat Am Enfermagem. 2012;20:35-43. <https://doi.org/10.1590/s0104-11692012000100006>.
- 536 23. Chen IH, Wang CH, Wang SY, Cheng SY, Yu TJ, Kuo SF. Mediating effects of shoulder-
537 arm exercise on the postoperative severity of symptoms and quality of life of women with
538 breast cancer. BMC Womens Health. 2020;20:101. [https://doi.org/10.1186/s12905-020-00968-](https://doi.org/10.1186/s12905-020-00968-w)
539 w.
- 540 24. Yuan R, Wei X, Ye Y, et al. The effects of the mirror therapy on shoulder function in patients
541 with breast cancer following surgery: a randomized controlled trial. J Cancer Surviv. 2023;
542 Online ahead of print. <https://doi.org/10.1007/s11764-023-01398-x>.
- 543 25. Warehime S, Dinkel D, Alonso W, Pozehl B. Long-term exercise adherence in patients with
544 heart failure: A qualitative study. Heart Lung. 2020;49:696-701.
545 <https://doi.org/10.1016/j.hrtlng.2020.08.016>.
- 546 26. Arinaga Y, Piller N, Sato F, et al. The 10-Min Holistic Self-Care for Patients with Breast
547 Cancer-Related Lymphedema: Pilot Randomized Controlled Study. Tohoku J Exp Med.
548 2019;247:139-147. <https://doi.org/10.1620/tjem.247.139>.
- 549 27. Jeffs E, Wiseman T. Randomised controlled trial to determine the benefit of daily home-
550 based exercise in addition to self-care in the management of breast cancer-related
551 lymphoedema: a feasibility study. Support Care Cancer. 2013;21:1013-1023.
552 <https://doi.org/10.1007/s00520-012-1621-6>.
- 553 28. Tsang S, Royse CF, Terkawi AS. Guidelines for developing, translating, and validating a
554 questionnaire in perioperative and pain medicine. Saudi J Anaesth. 2017;11:S80-S89.
555 https://doi.org/10.4103/sja.SJA_203_17.
- 556 29. Zhou K, Wang W, Li M, et al. Body image mediates the relationship between post-surgery

- 557 needs and health-related quality of life among women with breast cancer: a cross-sectional
558 study. *Health Qual Life Outcomes*. 2020;18:163. <https://doi.org/10.1186/s12955-020-01400-5>.
- 559 30. Chang L, Zhang S, Yan Z, Li C, Zhang Q, Li Y. Symptom burden, family resilience, and
560 functional exercise adherence among postoperative breast cancer patients. *Asia Pac J Oncol*
561 *Nurs*. 2022;9:100129. <https://doi.org/10.1016/j.apjon.2022.100129>.
- 562 31. Podsakoff PM, MacKenzie SB, Lee J-Y, Podsakoff NP. Common method biases in
563 behavioral research: a critical review of the literature and recommended remedies. *J Appl*
564 *Psychol*. 2003;88:879-903. <https://doi.org/10.1037/0021-9010.88.5.879>.
- 565 32. Altas EU, Demirdal Ü S. The effects of post-mastectomy lymphedema on balance,
566 Kinesiophobia and fear of falling. *J Community Health Nurs*. 2021;38:130-138.
567 <https://doi.org/10.1080/07370016.2021.1887564>.
- 568 33. Velthuis MJ, Peeters PH, Gijzen BC, et al. Role of fear of movement in cancer survivors
569 participating in a rehabilitation program: a longitudinal cohort study. *Arch Phys Med Rehabil*.
570 2012;93:332-338. <https://doi.org/10.1016/j.apmr.2011.08.014>.
- 571 34. Jiang W, Chen L. Analysis of the factors and moderating role of self-care ability among
572 patients with breast cancer-related lymphedema. *J Clin Nurs*. 2023;32:926-940.
573 <https://doi.org/10.1111/jocn.16495>.
- 574 35. Karaca-Mandic P, Solid CA, Armer JM, Skoracki R, Campione E, Rockson SG.
575 Lymphedema self-care: economic cost savings and opportunities to improve adherence. *Cost*
576 *Eff Resour Alloc*. 2023;21:47. <https://doi.org/10.1186/s12962-023-00455-7>.
- 577 36. Tsuchiya M, Masujima M, Kato T, et al. Knowledge, fatigue, and cognitive factors as
578 predictors of lymphoedema risk-reduction behaviours in women with cancer. *Support Care*

- 579 Cancer. 2019;27:547-555. <https://doi.org/10.1007/s00520-018-4349-0>.
- 580 37. Shahsavari H, Matory P, Zare Z, Taleghani F, Kaji MA. Effect of self-care education on the
581 quality of life in patients with breast cancer. *J Educ Health Promot.* 2015;4:70.
582 <https://doi.org/10.4103/2277-9531.171782>.
- 583 38. Mokhtari-Hessari P, Montazeri A. Health-related quality of life in breast cancer patients:
584 review of reviews from 2008 to 2018. *Health Qual Life Outcomes.* 2020;18:338.
585 <https://doi.org/10.1186/s12955-020-01591-x>.
- 586 39. Kim S, Han J, Lee MY, Jang MK. The experience of cancer-related fatigue, exercise and
587 exercise adherence among women breast cancer survivors: Insights from focus group
588 interviews. *J Clin Nurs.* 2020;29:758-769. <https://doi.org/10.1111/jocn.15114>.
- 589 40. Cormie P, Pumpa K, Galvão DA, et al. Is it safe and efficacious for women with
590 lymphedema secondary to breast cancer to lift heavy weights during exercise: a randomised
591 controlled trial. *J Cancer Surviv.* 2013;7:413-424. <https://doi.org/10.1007/s11764-013-0284-8>.
- 592 41. Li F, Liu W, Huo F, et al. Effect of Self-Controlled Exercise on Antioxidant Activity of Red
593 Blood Cells and Functional Recovery of Limbs in Patients with Breast Cancer after
594 Rehabilitation. *Iran J Public Health.* 2021;50:306-314.
595 <https://doi.org/10.18502/ijph.v50i2.5345>.
- 596 42. Kim M, So WY, Kim J. Relationships between Exercise Modality and Activity Restriction,
597 Quality of Life, and Hematopoietic Profile in Korean Breast Cancer Survivors. *Int J Environ
598 Res Public Health.* 2020;17:6899. <https://doi.org/10.3390/ijerph17186899>.

599 **Figure legends**

600 **Fig. 1.** Hypothesized model.

601 **Fig. 2.** Flow diagram of study recruitment process.

602 **Fig. 3.** The chain-mediating model of self-care and functional exercise adherence in the

603 correlation between kinesiophobia and QoL. ** $p < 0.01$, *** $p < 0.001$ was considered

604 statistically significant.

Journal Pre-proof

Table 1. Participants' characteristics and different variable scores (N = 274).

Variables	n (%)	Kinesiophobia score			Self-care score			PFEA score			QoL score		
		Mean	SD	Significant	Mean	SD	Significant	Mean	SD	Significant	Mean	SD	Significant
Age													
< 50 years	155 (56.6)	28.77	4.39	0.038*	41.78	5.02	0.002**	44.32	7.19	0.001**	81.32	12.19	0.004**
≥ 50 years	119 (43.4)	29.91	4.55		39.86	4.85		41.49	7.12		77.17	11.21	
BMI													
< 23 kg/m ²	92 (33.6)	28.99	4.33	0.468	40.98	5.06	0.918	42.48	7.57	0.323	79.20	12.29	0.751
≥ 23 kg/m ²	182 (66.4)	29.41	4.57		40.92	5.03		43.40	7.14		79.68	11.77	
Education level													
< senior high school	177 (64.6)	29.41	4.34	0.396	40.55	4.92	0.082	42.97	7.39	0.702	79.06	11.81	0.394
≥ senior high school	97 (36.4)	29.01	4.76		41.66	5.18		42.32	7.13		80.35	12.17	
Marital status													
Married/Partner	231 (84.3)	28.95	4.39	0.007**	41.05	5.00	0.436	43.30	7.30	0.266	80.17	11.91	0.035*
Single	43 (15.7)	30.95	4.65		40.40	5.21		41.96	7.19		76.00	11.56	
Monthly income													
< 2000 yuan	89 (32.5)	29.28	4.59	0.994	40.47	5.23	0.260	42.10	7.27	0.173	78.83	12.83	0.654
2000-5000 yuan	166 (60.6)	29.27	4.47		41.01	4.96		43.38	7.37		79.66	11.61	
> 5000 yuan	19 (6.9)	29.16	4.32		42.53	4.57		45.21	6.17		81.53	10.61	

Have children													
Yes	238 (86.9)	29.16	4.56	0.331	41.17	4.96	0.059	43.05	7.24	0.812	79.74	12.03	0.431
No	36 (13.1)	29.94	3.94		39.47	5.28		43.36	7.71		78.06	11.28	
Employment													
Unemployed	153 (55.8)	29.14	4.60	0.612	41.38	4.91	0.109	43.52	7.24	0.271	80.25	12.70	0.255
Employed	121 (44.2)	29.42	4.34		40.40	5.14		42.55	7.33		78.60	10.86	
BC duration													
< 4 years	123 (44.9)	28.54	4.71	0.015*	41.91	5.43	0.004**	43.72	7.49	0.195	81.36	12.01	0.021*
≥ 4 years	151 (55.1)	29.86	4.22		40.16	4.55		42.58	7.10		78.02	11.69	
Chemotherapy													
Yes	153 (55.8)	29.90	4.74	0.008**	40.56	4.85	0.157	42.12	6.99	0.013**	77.77	12.13	0.006**
No	121	28.46	4.02		41.43	5.23		44.32	7.50		81.73	11.34	
Radiotherapy													
Yes	138 (50.4)	30.01	4.46	0.006**	40.24	5.13	0.019*	42.33	7.38	0.080	76.80	10.56	<0.001***
No	136	28.51	4.40		41.66	4.84		43.87	7.13		82.27	12.63	
Tumor location													
Left	143 (52.2)	28.78	4.01	0.115	40.76	4.90	0.066	43.15	6.82	0.131	80.10	11.68	0.068
Right	126 (46.0)	29.72	4.93		41.34	5.14		43.28	7.83		79.33	12.16	
Bilateral	5 (1.8)	31.60	4.56		36.20	3.56		36.60	2.51		67.60	7.64	

BC stages													
Stage I-II	212 (77.4)	28.84	4.48	0.003**	41.24	5.15	0.077	43.57	7.40	0.044*	80.51	11.95	0.010*
Stage III-IV	62 (22.6)	30.73	4.24		39.95	4.47		41.45	6.70		76.11	11.31	
Surgery type													
MRM	171 (62.4)	29.16	4.41	0.714	41.28	5.00	0.364	43.60	7.34	0.288	80.20	11.98	0.209
BCS	84 (30.7)	29.30	4.58		40.37	5.17		42.42	7.16		79.11	12.06	
Others ^b	19 (6.9)	30.05	4.88		40.47	4.64		41.47	7.29		75.21	10.42	
Comorbidities ^c													
Yes	195 (71.2)	29.62	4.49	0.040*	40.71	5.10	0.230	42.67	7.12	0.130	78.59	11.70	0.042*
No	79 (28.8)	28.39	4.39		41.52	4.83		44.14	7.63		81.82	12.25	

Note. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Abbreviations: BC, breast cancer; BMI, Body Mass Index; BCS, breast conserving surgery; MRM, modified radical mastectomy; PFEA, postoperative functional exercise adherence; QoL, quality of life; SD, Standard deviation.

^aOne-way ANOVA test/independent t -test.

^bLumpectomy and axillary dissection or total mastectomy.

^cHypertension, diabetes, thyroid, osteoporosis, or dyslipidemia.

Table 2. Correlation between kinesiophobia, self-care, functional exercise adherence, and QoL in patients with BCRL (r).

Variables	Mean	SD	1.	2.	3.	4.	5.	6.	7.	8.	9.
1. Kinesiophobia	29.27	4.49	1								
2. Self-care	40.95	5.03	-0.481***	1							
3. PFEA	43.09	7.29	-0.564***	0.614***	1						
4. QoL	79.52	11.93	-0.680***	0.569***	0.713***	1					
5. PWB	14.82	2.69	-0.194***	0.198***	0.230***	0.464***	1				
6. EWB	13.80	2.96	-0.520***	0.366***	0.508***	0.749***	0.241***	1			
7. SWB	15.31	3.49	-0.441***	0.476***	0.492***	0.696***	0.155***	0.383***	1		
8. FWB	13.46	3.01	-0.355***	0.317***	0.341***	0.586***	0.700***	0.379***	0.342***	1	
9. BCS	18.78	4.88	-0.569***	0.447***	0.629***	0.802***	0.176***	0.538***	0.372***	0.311**	1

Note. *** $p < 0.001$. Abbreviations: BCRL, breast cancer related lymphedema; BCS, Breast Cancer Subscale for additional concerns; EWB, emotional well-being; FWB, functional well-being; PFEA, postoperative functional exercise adherence; PWB, physical well-being; QoL, quality of life; SD, standard deviation; SWB, social/family well-being.

Table 3. Regression model of the relationship between factors (N = 274).


Variables	QoL		Self-care		PFEA		QoL	
	β (95%CI)	<i>t</i>	β (95%CI)	<i>t</i>	β (95%CI)	<i>t</i>	β (95%CI)	<i>t</i>
Age	-0.084 (-0.233, 0.004)	-1.896	-0.115 (-0.126, -0.006)	-2.160*	-0.063 (-0.126, 0.022)	-1.394	-0.032 (-0.132, 0.069)	-0.619
Marital status	-0.019 (-3.507, 2.294)	-0.412	0.040 (-0.911, 2.021)	0.746	0.010 (-0.118, 0.624)	1.341	-0.035 (-3.558, 1.292)	-0.920
BC duration	-0.031 (-0.806, 0.389)	-0.689	-0.001 (-0.651, -0.047)	-2.276*	0.062 (-1.596, 1.977)	0.210	-0.020 (-0.640, 0.371)	-0.523
BC stage	-0.024 (-3.232, 1.885)	-0.518	-0.001 (-1.307, 1.279)	-0.021	-0.013 (-1.796, 1.352)	-0.278	-0.018 (-2.648, 1.626)	-0.471
Chemotherapy	0.038 (-1.242, 3.045)	0.828	0.002 (-1.064, 1.103)	0.035	0.053 (-0.536, 2.102)	1.169	0.015 (-1.446, 2.143)	0.383
Radiotherapy	0.113 (0.567, 4.808)	2.395*	0.067 (-0.395, 1.749)	1.243	-0.031 (-1.760, 0.857)	-0.680	0.106 (0.739, 4.294)	2.787**
Comorbidities	0.030 (-1.511, 3.107)	0.680	0.004 (-1.127, 1.207)	0.068	0.011 (-1.250, 1.591)	0.236	0.025 (-1.277, 2.580)	0.666
Kinesiophobia	-0.630 (-1.923, -1.427)	-13.276***	-0.441 (-0.620, -0.369)	-7.751***	-0.349 (-0.736, -0.398)	-6.598***	-0.351 (-1.181, -0.685)	-7.407***
Self-care					0.446 (0.500, 0.794)	8.650***	0.113 (0.041, 0.493)	2.322*
PFEA							0.420 (0.524, 0.853)	8.237***
R ²	0.488		0.264		0.483		0.646	
F	31.5845***		11.895***		27.355***		37.946***	

Note. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. Abbreviations: BC, breast cancer; CI, confidence interval; PFEA, postoperative functional exercise adherence; QoL, quality of life.

Table 4. Bootstrap analysis of the mediating effect of kinesiophobia and QoL (N = 274).

Paths	Effect	BootSE	BootLLCI	BootULCI	Effect ratio
Total effect	-1.675	0.126	-1.923	-1.427	
Direct effect	-0.933	0.126	-1.181	-0.685	55.7%
Total indirect effect	-0.742	0.082	-0.915	-0.592	44.3%
Ind1: Kinesiophobia → Self-care → QoL	-0.132	0.065	-0.267	-0.007	7.9%
Ind2: Kinesiophobia → PFEA → QoL	-0.390	-0.064	-0.516	-0.269	23.3%
Ind 3: Kinesiophobia → Self-care → PFEA → QoL	-0.220	-0.046	-0.320	-0.142	13.1%
Comparison1 (Indirect1 - Indirect2)	0.258	0.104	0.053	0.457	
Comparison2 (Indirect1 - Indirect3)	0.088	0.088	-0.081	0.273	
Comparison3 (Indirect2 - Indirect3)	-0.170	0.077	-0.313	-0.009	

Note. Abbreviations: CI, confidence interval; PFEA, postoperative functional exercise adherence; QoL, quality of life.



Journal Pre-proof

Flow Diagram




Fig. 2. Flow diagram of study recruitment process.

